

A restricted spectrum of *NRAS* mutations causes Noonan syndrome

Ion C Cirstea¹, Kerstin Kutsche², Radovan Dvorsky¹, Lothar Gremer¹, Claudio Carta³, Denise Horn⁴, Amy E Roberts^{5,6}, Francesca Lepri⁷, Torsten Merbitz-Zahradnik¹, Rainer König⁸, Christian P Kratz^{9,10}, Francesca Pantaleoni³, Maria L Dentici⁷, Victoria A Joshi^{5,11,12}, Raju S Kucherlapati⁵, Laura Mazzanti¹³, Stefan Mundlos⁴, Michael A Patton¹⁴, Margherita Cirillo Silengo¹⁵, Cesare Rossi¹⁶, Giuseppe Zampino¹⁷, Cristina Digilio¹⁸, Liborio Stuppia¹⁹, Eva Seemanova²⁰, Len A Pennacchio^{21,22}, Bruce D Gelb^{23–25}, Bruno Dallapiccola⁷, Alfred Wittinghofer²⁶, Mohammad R Ahmadian^{1,29}, Marco Tartaglia^{3,29} & Martin Zenker^{26–29}

Noonan syndrome, a developmental disorder characterized by congenital heart defects, reduced growth, facial dysmorphism and variable cognitive deficits, is caused by constitutional dysregulation of the RAS-MAPK signaling pathway. Here we report that germline *NRAS* mutations conferring enhanced stimulus-dependent MAPK activation account for some cases of this disorder. These findings provide evidence for an obligate dependency on proper *NRAS* function in human development and growth.

The *KRAS*, *NRAS* and *HRAS* genes encode highly conserved small GTPases that act as membrane-associated molecular switches controlling intracellular signal flow by cycling between active, GTP-bound states and inactive, GDP-bound states¹. Following the transmission of signals elicited by cell surface receptors, RAS proteins are activated by guanine nucleotide exchange factors. In the GTP-bound state, two regions, switch I and switch II, undergo a conformational change that enables

RAS proteins to bind to and activate effector proteins, including RAF and phosphatidylinositol-3-kinases². This interaction is terminated by hydrolysis of GTP to GDP, which promotes a switch toward the inactive RAS conformation. The low intrinsic GTPase activity of RAS proteins is enhanced by GTPase activating proteins (GAPs)³. Because of its nodal role in signal transduction, RAS function is required for many cellular processes, and its upregulation has been causally linked to oncogenesis and developmental disorders^{2,4}. These developmental disorders comprise a group of clinically related conditions characterized by facial dysmorphism, congenital cardiac defects, reduced growth and variable cognitive impairment and tumor predisposition. In these disorders, RAS signaling dysregulation is caused by germline mutations in *HRAS* or *KRAS* as well as in genes encoding modulators of RAS function (*PTPN11*, *SOS1*, *SHOC2*, *NF1* and *SPRED1*) or downstream signal transducers (*RAF1*, *BRAF*, *MEK1* and *MEK2*)^{4–7}. The common biological consequence of mutations associated with this group of diseases is altered, usually increased, signal traffic through the MAPK cascade.

Based on the evidence that Noonan syndrome (MIM163950) and the clinically related cardio-facio-cutaneous syndrome (MIM115150) are genetically heterogeneous, with mutations in known disease genes accounting for 70–80% of cases of each^{6,7}, a systematic scanning of the *NRAS* gene was performed on a cohort of 917 individuals negative for previously known mutations with a phenotype fitting or suggestive of these disorders (**Supplementary Methods**). We identified four unrelated individuals in this cohort heterozygous for the nucleotide changes C149T or G179A (resulting in the amino acid substitutions T50I and G60E, respectively) (**Fig. 1** and **Supplementary Fig. 1**). Three were sporadic cases, and parental DNA genotyping demonstrated the *de novo* origin of the mutations. The defects were documented in hair bulbs, urine exfoliated cells and/or buccal epithelial cell specimens, thus excluding a somatic event restricted to blood cells as the cause of the mutations (**Supplementary Fig. 1** and data not shown). In the family transmitting the trait, the mutation segregated with disease. All *NRAS* mutation-positive subjects showed typical clinical features of Noonan syndrome (**Supplementary Table 1** and **Supplementary Fig. 2**).

¹Institute of Biochemistry and Molecular Biology II, Heinrich-Heine University Medical Center, Düsseldorf, Germany. ²Institut für Humangenetik, Universitätsklinikum Hamburg-Eppendorf, Hamburg, Germany. ³Dipartimento di Ematologia, Oncologia e Medicina Molecolare, Istituto Superiore di Sanità, Rome, Italy. ⁴Institute of Medical Genetics, Charité, University Medicine of Berlin, Berlin, Germany. ⁵Harvard Medical School, Boston, Massachusetts, USA. ⁶Department of Cardiology, Children's Hospital Boston, Boston, Massachusetts, USA. ⁷Istituto di Ricovero e Cura a Carattere Scientifico-Casa Sollievo della Sofferenza, San Giovanni Rotondo and Casa Sollievo della Sofferenza-Istituto Mendel, Rome, Italy. ⁸Institute of Human Genetics, University of Frankfurt, Frankfurt, Germany. ⁹Clinical Genetics Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, Maryland, USA. ¹⁰Department of Paediatrics and Adolescent Medicine, University of Freiburg, Freiburg, Germany. ¹¹Partners HealthCare Center for Personalized Genetic Medicine, Boston, Massachusetts, USA. ¹²Department of Pathology, Massachusetts General Hospital, Boston, Massachusetts, USA. ¹³Dipartimento di Pediatria, Università degli Studi di Bologna, Bologna, Italy. ¹⁴Department of Clinical Genetics, St. Georges Hospital, London, UK. ¹⁵Dipartimento di Pediatria, Università di Torino, Turin, Italy. ¹⁶Unità Operativa di Genetica Medica, Policlinico S. Orsola-Malpighi, Bologna, Italy. ¹⁷Istituto di Clinica Pediatrica, Università Cattolica del Sacro Cuore, Rome, Italy. ¹⁸Sezione di Genetica Medica, Ospedale 'Bambino Gesù', Rome, Italy. ¹⁹Dipartimento di Scienze Biomediche, Università degli Studi 'G. d'Annunzio', Chieti, Italy. ²⁰Institute of Biology and Medical Genetics, Charles University, University Hospital Prague, Prague, Czech Republic. ²¹Genomics Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA. ²²US Department of Energy Joint Genome Institute, Walnut Creek, California, USA. ²³Center for Molecular Cardiology, ²⁴Departments of Pediatrics and ²⁵Genetics and Genomic Sciences, Mount Sinai School of Medicine, New York, New York, USA. ²⁶Max Planck Institute of Molecular Physiology, Department of Structural Biology, Dortmund, Germany. ²⁷Institute of Human Genetics, University Hospital of Erlangen, University of Erlangen-Nuremberg, Erlangen, Germany. ²⁸Institute of Human Genetics, University Hospital of Magdeburg, Otto von Guericke University, Magdeburg, Germany. ²⁹These authors contributed equally to this work. Correspondence should be addressed to M.Z. (martin.zenker@med.ovgu.de), M.T. (mtartaglia@iss.it) or R.A. (reza.ahmadian@uni-duesseldorf.de).

Received 22 June; accepted 19 October; published online 6 December 2009; doi:10.1038/ng.497




Figure 1 *NRAS* genomic organization and protein structure and position of Noonan syndrome-causing *NRAS* mutations. (a) Exon-intron structure of the human *NRAS* gene showing untranslated regions as gray boxes and coding exons as blue boxes. (b) Motifs and secondary structures of the *NRAS* protein. Functional motifs including the P-loop, switch I (Sw I), switch II (Sw II) and the hypervariable region (HVR) are highlighted. Underneath, secondary structural elements are shown as green arrows and blue cylinders representing β -sheets and α -helices, respectively. (c) Partial amino acid sequence alignment of human *NRAS*, *KRAS* and *HRAS* showing conservation of Thr50 and Gly60 (red). Orange boxes on top of the alignment mark amino acids comprising the switch I and II regions, and the yellow box marks the four amino acids of the β 2– β 3 loop. Red dotted lines in panels a to c indicate the positions of nucleotides and amino acids, respectively, affected by Noonan syndrome-causing mutations.


Both *NRAS* mutations affected amino acid residues highly conserved among RAS orthologs and paralogs (Fig. 1c and data not shown). Gly60 is located in the switch II region. A germline *KRAS* mutation affecting Gly60 (G60R) had previously been reported in cardio-facio-cutaneous syndrome⁸, and the somatic *NRAS* mutation G60E rarely occurs in malignancies (COSMIC database; see URLs below). Alterations of the neighboring Gln61 are frequent among oncogenic RAS alterations and are known to confer impaired GTPase function. In contrast, no somatic or germline RAS gene mutation affecting Thr50 has been described so far. Thr50 is an exposed residue located in the β 2– β 3 loop connecting the two switch regions (Fig. 1b) but is not predicted to be directly involved in GTP and GDP binding, GTP hydrolysis, or effector interaction (Supplementary Fig. 3).

To elucidate the consequences of the Noonan syndrome-causing *NRAS* mutations on protein function and intracellular signaling, we expressed each mutant as yellow fluorescent protein–*NRAS* fusion proteins in cells from the COS-7 line, and their functional properties *in vitro* were compared with wild-type *NRAS* and the oncogenic *NRAS* substitution G12V, where the latter is known to be GAP resistant and to promote constitutive activation of MAPK signaling. Expression of the *NRAS* substitutions T50I or G60E resulted in enhanced phosphorylation of MEK and ERK in the presence of serum or after epidermal growth factor (EGF) stimulation but, in contrast with the G12V *NRAS* substitution, had only marginal effects on signaling in starved cells (Fig. 2a–c). Of note, the G60E *NRAS* mutant, much like G12V,

accumulated constitutively in the active, GTP-bound form and appeared resistant to GAP stimulation. In contrast, the T50I *NRAS* mutant was not enriched in the GTP-bound state (Fig. 2d,e), indicating a different mechanism of functional dysregulation. Biochemical characterization of this alteration provided evidence that the T50I substitution does not substantially affect intrinsic GTPase activity, GAP association or GAP-stimulated hydrolysis. Similarly, no change was observed in guanine nucleotide exchange factor–catalyzed nucleotide release, effector association or allosteric activation of SOS1 (Supplementary Fig. 4).

To explore further the structural and functional consequences of the T50I change, we determined the crystal structure of the RAS mutant bound to Gpp(NH)p (an analog of GTP), but no substantial structural rearrangements were observed (Supplementary Table 2 and Supplementary Fig. 5). Molecular dynamics simulations of wild-type RAS indicated that Thr50 directly interacts with the polar heads of membrane phospholipids and is an integral part of a recently described functional region that controls RAS membrane orientation and signal output (Supplementary Fig. 6)⁹. According to this model, aside from the conformational changes involving the switch I and II regions, GTP loading of membrane-anchored RAS is predicted to rearrange a distinct region of the GTPase (comprising the β 2– β 3 loop and helix α 5) (Fig. 1c,d). This, in turn, would lead to a differential engagement of the C-terminal hypervariable region and helix α 4 with the membrane, ultimately altering the orientation of the entire G domain of RAS with respect to the membrane. Residues Asp47 and Glu49 have been shown to be directly involved in

Figure 2 Biochemical characterization of the Noonan syndrome-causing T50I and G60E *NRAS* mutants. (a,b) Determination of MEK, ERK and AKT phosphorylation levels in transiently transfected COS-7 cells cultured in medium with serum (a) or in basal medium (b). Expression of each of the three *NRAS* mutants (G12V, G60E and T50I) resulted in enhanced MEK and ERK phosphorylation in the presence of serum, whereas only the oncogenic G12V substitution substantially stimulated phosphorylation of MEK and ERK in serum-deprived cells. Contrasting with what was observed for the G12V change, no effect on AKT phosphorylation was detectable in cells expressing the Noonan syndrome-causing T50I or G60E mutant under both culture conditions. Asterisk beside the pAKT blot marks artifactual bands (Supplementary Fig. 7). Total amounts of MEK, ERK and AKT in cell lysates are shown for equal protein expression and loading. (c) EGF stimulation of serum-starved cells reproduced the effects on ERK phosphorylation observed in the presence of serum. (d) Determination of GTP-bound *NRAS* levels in COS-7 cells transiently expressing wild-type *NRAS* or each mutant in the presence of serum. Similar to the oncogenic G12V mutant, G60E, but not T50I, accumulated in the GTP-bound form. (e) The same experiment was performed under serum-free conditions. Both the G12V and G60E mutants were predominantly present in the active GTP-bound form, whereas the T50I mutant was not. Addition of purified RasGAP left GTP-bound levels of the G12V and G60E mutants unaffected, demonstrating GAP resistance of these mutants, in contrast to the T50I mutant. Total amounts of exogenous (yellow fluorescent protein-tagged) *NRAS* in cell lysates are shown for each experiment (a–e). Specificity of the antibody employed in each experiment is indicated below each panel. Wt, wild type.



the intermolecular binding network controlling RAS membrane orientation, and their substitution was found to enhance downstream signaling through GTP-bound RAS, presumably by keeping the GTPase in a more signaling-competent orientation⁹. Consistent with this model, the RAS-binding domain of RAF1 is able to recognize the relative membrane orientation of GTP-bound RAS, which is functionally translated into different signaling outputs⁹. Based on this model, we hypothesize that the T50I substitution might alter RAS orientation in a fashion similar to that documented for the D47A and E49A substitutions⁹ (Supplementary Fig. 6), bolstering the interaction of GTP-bound RAS with its effectors. This proposed mechanism would explain the enhanced signal flow through the MAPK cascade in the absence of any substantial accumulation of the T50I NRAS substitution in its GTP-bound form (Fig. 2). Taken together, these findings support the idea that Noonan syndrome-causing NRAS mutants are weak hypermorphs. Although the G60E NRAS substitution accumulates in its GTP-bound form to mimic the consequences of oncogenic substitutions affecting Gly12 or Gly13, the weakened effector binding of this mutant might account for its damped downstream effect compared to the oncogenic G12V alteration.

Here we reported that Noonan syndrome can be caused by altered signaling through NRAS. Similar to what has been observed for germline KRAS mutations causing Noonan syndrome and cardio-facio-cutaneous syndrome, germline NRAS alterations do not affect residues commonly mutated in human cancers and appear to be less activating in dysregulating intracellular signaling *in vitro* compared with cancer-associated somatic mutations. Considering the large size of the cohorts of affected individuals included in this study, germline NRAS mutations are estimated to be substantially less frequent than those affecting KRAS or HRAS⁷. Although we favor the hypothesis that NRAS mutations rarely occur as germline defects because of their severe consequences on embryonic and fetal development, we cannot exclude the possibility that mutations other than those described here may cause phenotypes not recognized as Noonan syndrome or related disorders. Notably, Oliveira and colleagues reported on an individual with a germline NRAS missense change (G13D) and an autoimmune lymphoproliferative disorder, without any signs of Noonan syndrome¹⁰. Although their observation has not been replicated so far, their findings could be consistent with the occurrence of non-overlapping allele-specific phenotypic effects resulting from different NRAS mutations than those described here.

The identification of activating NRAS mutations in Noonan syndrome provides evidence for an obligate dependency on proper function of this small GTPase in human development and growth. Even though functional redundancy among RAS proteins does exist, biological differences in these proteins are also well documented^{11,12}. Recent evidence gathered from mouse models demonstrated that these GTPases play distinct roles in development. Although loss of Kras isoforms is lethal at the embryonic stage, disruption of mouse *Hras* and *Nras* does not meaningfully perturb development. Consistently, it has been documented that RAS proteins generate distinct signal outputs despite their interaction with a shared set of activators and effectors^{11,12}, and preferential associations between specific human cancers and individual RAS genes have also been established (see COSMIC database). These and recent findings regarding Noonan syndrome and related disorders extend this concept to germline mutations and human development, showing that distinct clusters of activating mutations in the three RAS genes underlie a group of clinically related mendelian traits. Notably, whereas a small number of mutations in HRAS and NRAS underlie, respectively, Costello syndrome (MIM218040) and Noonan syndrome, a more diverse set of missense

changes in KRAS results in a markedly broad phenotypic continuum¹³. Further studies are required to understand such genotype-phenotype correlations. The localization of RAS isoforms to distinct signaling nanoclusters and their ability to differentially activate canonical and non-canonical pathways, rather than distinct tissue- or cell-specific expression patterns, are believed to underlie the functional diversity of these GTPases^{11,12,14}. Accordingly, phenotypic differences between Noonan, cardio-facio-cutaneous and Costello syndromes might be ascribed to different developmental consequences strictly depending upon these mechanisms, although the obvious similarities of these disorders testify that there are important functions in developmental programs that are shared by all members of this family of GTPases.

This study was approved by the Institutional Review Boards at the University of Erlangen-Nuremberg, the Istituto Superiore di Sanità, IRCCS-Casa Sollievo della Sofferenza, and the Children's Hospital Boston and Partners HealthCare, Boston. Informed consent was obtained from all subjects participating in the study and/or their parents.

URLs. COSMIC database, <http://www.sanger.ac.uk/genetics/CGP/cosmic/>.

Note: Supplementary information is available on the Nature Genetics website.

ACKNOWLEDGMENTS

We are grateful to the subjects and their families for participating in this study. We thank A. Diem and B. Quinger (Institute of Human Genetics, Erlangen, Germany), T. Squatriti and S. Venanzi (Istituto Superiore di Sanità, Rome, Italy) and I. Jantke and S. Lorenz (Institut für Humangenetik, Hamburg, Germany) for skilful technical assistance. The X-ray diffraction datasets were collected at the Swiss Light Source, beamline X10SA, Paul Scherrer Institut, Villigen, Switzerland. This work was supported by grants from the European Research Area Network for research programs on rare diseases (E-Rare) 2009 to M.T., M.Z. and M.R.A. (European Network on Noonan Syndrome and Related Disorders), German Research Foundation (DFG) to M.R.A. (AH 92/5-1) and C.P.K. and M.Z. (KR 3473/1-1 and ZE 524/4-1), Nationales Genomforschungsnetz Plus program of the German Ministry of Science and Education to M.R.A. (BMBF grant 01GS08100), and Telethon-Italy (GGP07115), 'Associazione Italiana Sindromi di Costello e Cardiofaciocutanea' and 'Collaborazione Italia-USA/malattie rare' to M.T.

AUTHOR CONTRIBUTIONS

K.K., C.C., D.H., F.L., F.P., M.L.D., V.A.J. and C.R. performed mutation analysis and genotyping. L.A.P. performed high-throughput resequencing. L.S. performed the NRAS copy number analysis. R.K., A.E.R., L.M., S.M., M.A.P., M.C.S., G.Z., C.D., E.S., B.D. and M.Z. obtained DNA specimens from affected individuals and did clinical evaluations. R.D. and L.G. determined crystal structure. R.D. performed molecular dynamics simulations and structural analyses. I.C.C., L.G., T.M.-Z. and A.W. carried out biochemical and functional studies. C.P.K., R.S.K. and B.D.G. did project planning and data analysis. K.K. participated in preparation of the manuscript. M.R.A., M.T. and M.Z. carried out the project planning and participated in data analysis and preparation of the manuscript.

Published online at <http://www.nature.com/naturegenetics/>.

Reprints and permissions information is available online at <http://npg.nature.com/reprintsandpermissions/>.

- Vetter, I.R. & Wittinghofer, A. *Science* **294**, 1299–1304 (2001).
- Karnoub, A.E. & Weinberg, R.A. *Nat. Rev. Mol. Cell Biol.* **9**, 517–531 (2008).
- Ahmadian, M.R., Hoffmann, U., Goody, R.S. & Wittinghofer, A. *Biochemistry* **36**, 4535–4541 (1997).
- Schubert, S., Shannon, K. & Bollag, G. *Nat. Rev. Cancer* **7**, 295–308 (2007).
- Cordeddu, V. *et al. Nat. Genet.* **41**, 1022–1026 (2009).
- Denayer, E., de Ravel, T. & Legius, E. *J. Med. Genet.* **45**, 695–703 (2008).
- Tidyman, W.E. & Rauen, K.A. *Curr. Opin. Genet. Dev.* **19**, 230–236 (2009).
- Niihori, T. *et al. Nat. Genet.* **38**, 294–296 (2006).
- Abankwa, D. *et al. EMBO J.* **27**, 727–735 (2008).
- Oliveira, J.B. *et al. Proc. Natl. Acad. Sci. USA* **104**, 8953–8958 (2007).
- Hancock, J.F. *Nat. Rev. Mol. Cell Biol.* **4**, 373–384 (2003).
- Quinlan, M.P. & Settleman, J. *Future Oncol.* **5**, 105–116 (2009).
- Zenker, M. *et al. J. Med. Genet.* **44**, 131–135 (2007).
- Haigis, K.M. *et al. Nat. Genet.* **40**, 600–608 (2008).